

Technical Data Sheet

InVivoMAb anti-mouse IL-15



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0315
Clone: AIO.3
Isotype: Rat IgG2a, λ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Recombinant mouse IL-15
Reported Applications: *in vivo* IL-15 neutralization
in vitro IL-15 neutralization
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_2754553](https://abnova.com/AB_2754553)
Molecular Weight: 150 kDa

Description

The AIO.3 monoclonal antibody reacts with mouse IL-15. IL-15 is a pro-inflammatory cytokine that is produced mainly by dendritic cells, epithelial cells, fibroblasts, and monocytes. IL-15 plays important roles in the immune response and shares many functions with IL-2. IL-15 has been shown to stimulate the proliferation of activated T cells, NK cells, and B cells, and induce antibody production by B cells stimulated with anti-IgM or CD40L. In addition, IL-15 promotes the development of dendritic cells and induces the production of proinflammatory cytokines from macrophages. IL-15 has also been shown to play a role in several inflammatory disorders, including rheumatoid arthritis, psoriasis and pulmonary inflammatory diseases. Emerging data suggest that there is a beneficial effect of IL-15 neutralization in models of psoriasis and diabetes. The AIO.3 antibody has been shown to neutralize the bioactivity of IL-15 *in vitro* and *in vivo*.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0315?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC
<https://bioxcell.com>
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC